

***Bcl10* Cas9-KO Strategy**

Designer: Daohua Xu

Reviewer: Huimin Su

Design Date: 2019-8-14

Project Overview

Project Name

Bcl10

Project type


Cas9-KO

Strain background

C57BL/6JGpt

Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Bcl10* gene. The schematic diagram is as follows:



- The *Bcl10* gene has 3 transcripts. According to the structure of *Bcl10* gene, exon2 of *Bcl10*-201(ENSMUST00000029842.9) transcript is recommended as the knockout region. The region contains 289bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Bcl10* gene. The brief process is as follows: CRISPR/Cas9 system were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.

- According to the existing MGI data, about one-third of homozygous null embryos die exhibiting exencephaly. Surviving mutants display immunological defects including severe immunodeficiency, abnormal B cell development and function, and impaired humoral response to bacterial infection.
- The *Bcl10* gene is located on the Chr3. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)

Bcl10 B cell leukemia/lymphoma 10 [Mus musculus (house mouse)]

Gene ID: 12042, updated on 13-Mar-2020

Summary

Official Symbol Bcl10 provided by [MGI](#)

Official Full Name B cell leukemia/lymphoma 10 provided by [MGI](#)

Primary source [MGI:MGI:1337994](#)

See related [Ensembl:ENSMUSG00000028191](#)

Gene type protein coding

RefSeq status VALIDATED

Organism [Mus musculus](#)

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as AI132454, BCL-10, C81403, CARMEN, CIPER, CLAP, ME10, cE10

Expression Ubiquitous expression in large intestine adult (RPKM 10.2), bladder adult (RPKM 8.8) and 28 other tissues [See more](#)


Orthologs [human](#) [all](#)

Transcript information (Ensembl)


The gene has 3 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Bcl10-201	ENSMUST00000029842.8	1949	233aa	Protein coding	CCDS17897	B7ZWE5 Q9Z0H7	TSL:1 GENCODE basic APPRIS P1
Bcl10-203	ENSMUST00000198122.1	660	No protein	Processed transcript	-	-	TSL:3
Bcl10-202	ENSMUST00000197842.1	2368	No protein	Retained intron	-	-	TSL:NA


The strategy is based on the design of *Bcl10-201* transcript,the transcription is shown below:




Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, about one-third of homozygous null embryos die exhibiting exencephaly.

Surviving mutants display immunological defects including severe immunodeficiency, abnormal B cell development and function, and impaired humoral response to bacterial infection.

If you have any questions, you are welcome to inquire.
Tel: 400-9660890

