




## Genotyping Report

Strain ID	T052706	Strain Type	KI(Cas9)	Genetic Background	BALB/cJGpt
Designer	Dongdong Zhang	Gene Name			<i>Scnn1a-P2A-iCre</i>

### 1. Strategy of Genotyping



Wild type: ①PCR reaction obtains none band; ②PCR reaction obtains a WT band.

Heterozygote: ①PCR reaction obtains a Targeted band; ②PCR reaction obtains a WT band.


Homozygote: ①PCR reaction obtains a Targeted band; ②PCR reaction obtains none band.

Note: The sizes of WT and Targeted band are shown below. For ②PCR reaction, because the WT band is much smaller than the target band, it is likely to produce dominant amplification, the reaction is only used to judge whether there is a WT allele.

### 2. Primer Information

PCR No.	Primer No.	Primer Name	Sequence	Band Size
①5'arm	F1	XM004172-Scnn1a-wt-tF1	TCCACCTGCCTATGCTACCCCTA	WT:0bp Targeted:776bp
	R1	iCre-5tR1	TGCCAATGTGGATCAGCATT	
②WT	F2	XM004172-Scnn1a-wt-tF1	TCCACCTGCCTATGCTACCCCTA	WT:334bp Targeted:1462bp
	R2	XM004172-Scnn1a-wt-tR1	AAGTCTGGCTTGGCTGATCCAA	

### 3. Gel Image & Conclusion





Note: P:Positive control; WT: Wildtype control; B: Blank control (ddH<sub>2</sub>O); M: DNA Ladder

① Control (WT) : It is an important reference mark for whether the PCR reaction is successful and whether the product band position and size meet the theoretical requirements.

② Control (B) : PCR amplification was performed without template in the PCR reagent to monitor whether the reagent was contaminated.

#### 4. PCR Condition

PCR Reaction Component		
Seg.	reaction component	Volume (μl)
1	2 × Rapid Taq Master Mix (Vazyme P222)	12.5
2	ddH <sub>2</sub> O	9.5
3	Primer A(10pmol/μl)	1
4	Primer B(10pmol/μl)	1
5	Template(≈100ng/μl)	1

##### PCR program ① priority selection

Seg.	Temp.	Time	Cycle
1	95 °C	5min	
2	98 °C	30s	
3	65 °C * (-0.5 °C/cycle)	30s	
4	72 °C	45s*	
5	98 °C	30s	
6	55 °C *	30s	
7	72 °C	45s*	
8	72 °C	5min	
9	10 °C	hold	

##### PCR program ② the second choice

Seg.	Temp.	Time	Cycle
1	95 °C	5min	
2	98 °C	30s	
3	58 °C *	30s	
4	72 °C	45s*	
5	72 °C	5min	
6	10 °C	hold	

Note\*: Annealing temperature and extension time can be determined according to the actual amplification situation and amplification enzyme efficiency.



江苏集萃药康生物科技股份有限公司  
GemPharmatech Co.,Ltd