


Genotyping Report

Strain ID	T010975	Strain Type	CKO(Cas9)	Genetic Background	C57BL/6JGpt
Designer	Ninghui Zhao	Gene Name	<i>Irf7</i>		

1. Strategy of Genotyping




Wild type: ①PCR reaction obtains a single WT band; ②PCR reaction obtains a single WT band.
Heterozygote: ①PCR reaction obtains a WT band and a Targeted band; ②PCR reaction obtains a WT band and a Targeted band.
Homozygote: ①PCR reaction obtains a single Targeted band; ②PCR reaction obtains a single Targeted band.
Note: The sizes of WT and Targeted band are shown below.

2. Primer Information

PCR No.	Primer No.	Sequence	Band Size
①(5'arm)	T010975(P1)-F1	GCTAACCACAAGCAAGTCCCCAC	WT: 291bp Targeted: 396bp
	T010975(P1)-R1	CTTCCCTGGAAGTCACTGTGGAGA	
②(3'arm)	T010975(P1)-F2	AGGGAGGTACATGGCCTCTGTTCAT	WT: 305bp Targeted: 411bp
	T010975(P1)-R2	TTGCTGAGGCTCACTTCTTCCC	

3. Gel Image & Conclusion



Note: P: Positive control; WT: Wildtype control; B: Blank control (ddH₂O); M: DNA Ladder

① Control (WT) : It is an important reference mark for whether the PCR reaction is successful and whether the product band position and size meet the theoretical requirements.

② Control (B) : PCR amplification was performed without template in the PCR reagent to monitor whether the reagent



was contaminated.

4. PCR Condition

PCR Reaction Component			
Seg.	reaction component	Volume (μ l)	
1	2 \times Taq Master Mix , Dye Plus, (Vazyme P112-03)	12.5	
2	ddH ₂ O	9.5	
3	Primer A(10pmol/ μ l)	1	
4	Primer B(10pmol/ μ l)	1	
5	Template(\approx 100ng/ μ l)	1	
PCR program			
Seg.	Temp.	Time	Cycle
1	95°C	5min	
2	98°C	30s	20 \times
3	65°C (-0.5°C/cycle)	30s	
4	72°C	45s	
5	98°C	30s	
6	55°C	30s	20 \times
7	72°C	45s	
8	72°C	5min	
9	10°C	hold	