

		Genotyp	ing Report		-< 2
Strain ID	T052299	Strain Type	CKO(Cas9)	Genetic Background	C57BL/6JGpt
Designer	Ya'nan Xu	Gene Name	· · / ×	Ptpn18	6
and a		12hz		ans.	
Strategy of (	Genotyping	×,	3	3 <sup>6</sup>	7
	F1 → ①5'arm	R1 ◀	F2 (2)3	'arm R2	
/		Flox		//	
//					

1 2 PCR Reaction Number - Sequencing Region

Wild type: ①PCR reaction obtains a single WT band; ②PCR reaction obtains a single WT band. Heterozygote: ①PCR reaction obtains a WT band and a Targeted band; ②PCR reaction obtains a WT band and a Targeted band. Homozygote: ①PCR reaction obtains a single Targeted band; ②PCR reaction obtains a single Targeted band.

Note: The sizes of WT and Targeted band are shown below.

Primer

Loxp

## 2. Primer Information

PCR No.	Primer No.	No. Sequence		
(1)(5'arm)	T052299-F1	GCATAAACAGGAAGAAGGTGTCTC	WT: 337bp	
	T052299-R1	CAGCTTCAGGGATTCAATAGCCT	Targeted:442bp	
2)(3'arm)	T052299-F2	99-F2 GCTGTAACAGACAACATGACCATGC		
	T052299-R2 CCTCTCTGGCTTCTGCATACAGT		- Targeted:441bp	

## 3. Gel Image & Conclusion





**江苏集萃药康生物科技股份有限公司** GemPharmatech Co.,Ltd

① Control (WT) : It is an important reference mark for whether the PCR reaction is successful and whether the product band position and size meet the theoretical requirements.

② Control (B) : PCR amplification was performed without template in the PCR reagent to monitor whether the reagent was contaminated.

## 4. PCR Condition

	Contraction Contraction	Contra Maria		
4. PCR Condition		÷	and set	
PCR Reaction Component		2.	Mahuma (ul)	
Seg.	reaction component 2 × Rapid Taq Master Mix (Vazyme P222)		Volume (μl) 12.5	
1	ddH2O	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	9.5	
2			· 25 · · · · · · · · · · · · · · · · · ·	
3	Primer A(10pmol/µl)	×	1 %	
4	Primer B(10pmol/µl)	°C/	1 7	
5	Template(≈100ng/µl)		1 0	
PCR program ① pri	ority selection	s/x	$\gamma_{S,}$	
Seg.	Temp.	Time	Cycle	
1	95℃	5min	Mar	
2	98°C	30s	20×	
3	65℃*(-0.5℃/cycle)	30s	K. G.	
4 <u></u>	72°C	45s*	and it	
5	98°C	30s	20× 20×	
6	55°C*	30s	~ ~~~~	
7	72°C	45s*		
8	72°C	5min	B. CA	
9	10°C	hold		
PCR program $ extsf{@}$ th	e second choice	Co no	20	
Seg.	Temp.	Time	Cycle	
1 73%	95°C	5min	Marco Marco	
2	98°C	30s	35×	
3	58°C*	30s		
4 75,	72°C	45s*		
5	72°C	5min	na la	
6	10°C	hold	1 m 2	
6	10°C	hold	173 <sub>60</sub>	

Note\*: Annealing temperature and extension time can be determined according to the actual amplification situation and amplification enzyme efficiency.



