




## Genotyping Report

Strain ID	T006947	Strain Type	KI(Cas9)	Genetic Background	C57BL/6JGpt
Designer	Tianjiao Wang	Gene Name			<i>Clql2-iCre</i>

### 1. Strategy of Genotyping



Wild type: ①PCR reaction obtains none band; ②PCR reaction obtains a WT band.

Heterozygote: ①PCR reaction obtains a Targeted band; ②PCR reaction obtains a WT band.

Homozygote: ①PCR reaction obtains a Targeted band; ②PCR reaction obtains none band.


Note: The sizes of WT and Targeted band are shown below. For ②PCR reaction, because the WT band is much smaller than the target band, it is likely to produce dominant amplification, the reaction is only used to judge whether there is a WT allele.

### 2. Primer Information

PCR No.	Primer No.	Sequence	Band Size
①5'arm	T006947-F1	GGGCAGTCTGGTACTTCCAAGCT	WT:0bp Targeted:333bp
	T006947-R1	GGCCTCTGGGAAACAAGATG	
②WT	T006947-F2	CAGCAAAACCTGGCTGTGGATC	WT:412bp Targeted:0bp
	T006947-R2	ATGAGCCACCATGTGGGTGTC	

### 3. Gel Image & Conclusion





#### 4. PCR Condition

PCR Reaction Component			
Seg.	reaction component	Volume (μl)	
1	2 × Rapid Taq Master Mix (Vazyme P222)	12.5	
2	ddH <sub>2</sub> O	9.5	
3	Primer A(10pmol/μl)	1	
4	Primer B(10pmol/μl)	1	
5	Template(20~80ng/μl)	1	

#### PCR program I priority selection

Seg.	Temp.	Time	Cycle
1	95 °C	5min	20×
2	98 °C	30s	
3	65 °C * (-0.5 °C/cycle)	30s	
4	72 °C	45s*	
5	98 °C	30s	
6	55 °C *	30s	
7	72 °C	45s*	
8	72 °C	5min	
9	10 °C	hold	

#### PCR program II the second choice

Seg.	Temp.	Time	Cycle
1	95 °C	5min	
2	98 °C	30s	35×



江苏集萃药康生物科技股份有限公司  
GemPharmatech Co.,Ltd

3	58°C*	30s	
4	72°C	45s*	
5	72°C	5min	
6	10°C	hold	

Note\*: Annealing temperature and extension time can be determined according to the actual amplification situation and amplification enzyme efficiency.