

C57BL/6JGpt-H11-Vil1-iCre

Strain Name: C57BL/6JGpt-H11^{em1Cin(Vil1-icre)}/Gpt

Strain Type: Knock-in

Strain Number: T004714

Background: C57BL/6JGpt

Description

This mouse strain expresses codon optimized iCre recombinase ^[1] under the control of the 9 kb mouse *Vil1* promoter, the construct was precisely inserted into the H11 safe harbor site in mouse Chr11 by CRISPR/Cas9 technology. When crossed with a strain with loxP site flanked sequence in its genome, Cre-mediated recombination will result in excision of the DNA fragment between the two loxPs in epithelia of intestine and colon. Recombinase activity was also detected in a proportion of cells in stomach, kidney and liver. Note: mild sperm activity was detected by PCR detection of loxP recombination.

Strategy




Fig.1 Schematic diagram of C57BL/6JGpt-H11-Vil1-iCre model strategy.

Applications


1. Cre tool mice for specific induction of loxP recombination in intestinal epithelia of the duodenum, jejunum, ileum, and colon ^[2].

Data support

1. Validation methods & notes

H11-Vil1-iCre mice was crossed with CAG-loxp-ZsGreen-Stop-loxp-tdTomato mice with ubiquitous reporter expression (hereafter referred as CAG-G/R mice), Cre-mediated recombination will lead to excision of ZsGreen and the stop cassette and expression of tdTomato, thus loss of green fluorescence and gain of red fluorescence will indicate Cre activity. Fluorescence imaging of frozen sections were performed to exhibit Cre activity in various tissues and organs. Imaging of sections were performed under a 200x microscopy. Note: these results may only represent the activity of Cre in this strain at the identical stage. Recombinase activity may be different at other stages in your application.

2. Images of tissues and organs with obvious Cre activity







Fig 2. Fluorescence imaging of tissues and organs with obvious Cre activity.

Organ name was indicated in the left top of each subfigure group. Cre-: CAG-G/R single positive individuals; Cre+: H11-Vil1-iCre, CAG-G/R double positive individuals.

3. Images of tissues and organs with little or no Cre activity






Fig 3. Fluorescence imaging of tissues and organs with little or no Cre activity.

Organ name was indicated in the left top of each subfigure group. Cre-: CAG-G/R single positive individuals; Cre+: H11-Vil1-iCre, CAG-G/R double positive individuals.

Reference

1. Shimshek D R, Kim J, Hübner M R, et al. "Codon-improved Cre recombinase (iCre) expression in the mouse." *genesis* 2002, 32(1): 19-26.
2. Madison BB, Dunbar L, Qiao XT, et al. Cis elements of the villin gene control expression in restricted domains of the vertical (crypt) and horizontal (duodenum, cecum) axes of the intestine. *J Biol Chem.* 2002, 277(36):33275-83.