

Slc46a1 Cas9-KO Strategy

Designer: Xueting Zhang

Reviewer: Yanhua Shen

Date: 2020-02-12

Project Overview

Project Name

Slc46a1

Project type

Cas9-KO

Strain background

C57BL/6JGpt

Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Slc46a1* gene. The schematic diagram is as follows:



- The *Slc46a1* gene has 3 transcripts. According to the structure of *Slc46a1* gene, exon1-exon3 of *Slc46a1-201* (ENSMUST00000001126.3) transcript is recommended as the knockout region. The region contains start codon ATG. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Slc46a1* gene. The brief process is as follows: CRISPR/Cas9 system

- According to the existing MGI data, Mice homozygous for a knock-out allele exhibit increased circulating and liver levels of N-homocysteine and total homocysteine.
- Transcript *Slc46a1*-203 may not be affected.
- The *Slc46a1* gene is located on the Chr11. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)

Slc46a1 solute carrier family 46, member 1 [*Mus musculus* (house mouse)]

Gene ID: 52466, updated on 12-Aug-2019

Summary

- Official Symbol** Slc46a1 provided by [MGI](#)
- Official Full Name** solute carrier family 46, member 1 provided by [MGI](#)
- Primary source** [MGI:MGI:1098733](#)
- See related** [Ensembl:ENSMUSG00000020829](#)
- Gene type** protein coding
- RefSeq status** VALIDATED
- Organism** [Mus musculus](#)
- Lineage** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
- Also known as** HCP1; Pcft; D11Ert18e; 1110002C08Rik
- Expression** Biased expression in duodenum adult (RPKM 274.9), small intestine adult (RPKM 108.1) and 8 other tissues [See more](#)
- Orthologs** [human](#) [all](#)

Genomic context

Location: 11 B5; 11 46.74 cM See Slc46a1 in [Genome Data Viewer](#)

Exon count: 5


Annotation release	Status	Assembly	Chr	Location
108	current	GRCm38.p6 (GCF_000001635.26)	11	NC_000077.6 (78465701..78471945)
Build 37.2	previous assembly	MGSCv37 (GCF_000001635.18)	11	NC_000077.5 (78279203..78285447)

Transcript information (Ensembl)


The gene has 3 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Slc46a1-201	ENSMUST00000001126.3	1978	459aa	Protein coding	CCDS25104	Q6PEM8	TSL:1 GENCODE basic APPRIS P1
Slc46a1-202	ENSMUST00000146431.2	749	98aa	Nonsense mediated decay	-	F6QD87	TSL:3
Slc46a1-203	ENSMUST00000180786.1	1834	No protein	Retained intron	-	-	TSL:NA


The strategy is based on the design of *Slc46a1-201* transcript,The transcription is shown below




Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Mice homozygous for a knock-out allele exhibit increased circulating and liver levels of N-homocysteine and total homocysteine.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

