




## Genotyping Report

Strain ID	T013270	Strain Type	CKO(Cas9)	Genetic Background	C57BL/6JGpt
Designer	Ya'nan Xu	Gene Name			<i>Mmp16</i>

### 1. Strategy of Genotyping



Wild type: ①PCR reaction obtains a single WT band; ②PCR reaction obtains a single WT band.

Heterozygote: ①PCR reaction obtains a WT band and a Targeted band; ②PCR reaction obtains a WT band and a Targeted band.


Homozygote: ①PCR reaction obtains a single Targeted band; ②PCR reaction obtains a single Targeted band.


Note: The sizes of WT and Targeted band are shown below.

### 2. Primer Information

PCR No.	Primer No.	Primer Name	Sequence	Band Size
①(5'arm)	F1	JS04618-Mmp16-5wt-tF1	GCACATTCTGAGAATATCCAGCTACAG	WT:258bp Targeted:360bp
	R1	JS04618-Mmp16-5wt-tR1	AAGATTCCATGTGTTCTCCAGGAACC	
②(3'arm)	F2	JS04618-Mmp16-3wt-tF1	CCCAATAGTGAAAGTTCTCTAAATGGAA	WT:282bp Targeted:383bp
	R2	JS04618-Mmp16-3wt-tR1	AAAGTAATCGCCCCCAGTTGTAC	

### 3. Gel Image & Conclusion





#### 4. PCR Condition

##### PCR Reaction Component

Seg.	reaction component	Volume (μl)
1	2 × Rapid Taq Master Mix (Vazyme P222)	12.5
2	ddH <sub>2</sub> O	9.5
3	Primer A(10pmol/μl)	1
4	Primer B(10pmol/μl)	1
5	Template(20~80ng/μl)	1

##### PCR program I priority selection

Seg.	Temp.	Time	Cycle
1	95 °C	5min	
2	98 °C	30s	20×
3	65 °C * (-0.5 °C/cycle)	30s	
4	72 °C	45s*	
5	98 °C	30s	15×
6	55 °C *	30s	
7	72 °C	45s*	
8	72 °C	5min	
9	10 °C	hold	

##### PCR program II the second choice

Seg.	Temp.	Time	Cycle
1	95 °C	5min	
2	98 °C	30s	35×



江苏集萃药康生物科技股份有限公司  
GemPharmatech Co.,Ltd

3	58°C*	30s	
4	72°C	45s*	
5	72°C	5min	
6	10°C	hold	

Note\*: Annealing temperature and extension time can be determined according to the actual amplification situation and amplification enzyme efficiency.