




## Genotyping Report

Strain ID	T059535	Strain Type	KO(Cas9)	Genetic Background	C57BL/6JGpt
Designer	Zifan Lin	Gene Name			<i>Apc</i>

### 1. Strategy of Genotyping



Wild type: ①PCR reaction obtains a single WT band; ②PCR reaction obtains a single WT band.

Heterozygote: ①PCR reaction obtains a WT band and a KO band; ②PCR reaction obtains a WT band.

Homozygote: ①PCR reaction obtains a single KO band; ②PCR reaction without product.

Note: 1)The sizes of WT and Targeted band are shown below.


2) If the WT band is too large, it may not be possible to obtain a WT band.

### 2. Primer Information

PCR No.	Primer No.	Primer Name	Sequence	Band Size
PCR①	F1	T059535(P1)-F1	TCATTCTCCCCTACTTAGATGGCAT	WT: 12206bp KO: 626bp
	R1	T059535(P1)-R1	TGTGTGAGGGCATCAGATGTTG	
PCR②	F2	T059535(P1)-F2	CCTGCATTAGTACGCCAGTCTACTTT	WT: 325bp KO:0bp
	R2	T059535(P1)-R2	GGTTGATTGGTAGTCTGGAAGGACT	

### 3. Gel Image

gtgttcaactgagacagagacccgtact---11580bp---caaaggaaaggattgacatcagattccac



Note: P: Heterozygous samples; WT: Wildtype control; B: Blank control (ddH<sub>2</sub>O); M: DNA Ladder

① Control (WT) : It is an important reference mark for whether the PCR reaction is successful and whether the



product band position and size meet the theoretical requirements.

② Control (B) : PCR amplification was performed without template in the PCR reagent to monitor whether the reagent was contaminated.

#### 4. PCR Condition

PCR Reaction Component			
Seg.	reaction component	Volume (μl)	
1	2 × Rapid Taq Master Mix (Vazyme P222)	12.5	
2	ddH <sub>2</sub> O	9.5	
3	Primer A(10pmol/μl)	1	
4	Primer B(10pmol/μl)	1	
5	Template(20~80ng/μl)	1	
PCR program I (priority selection)			
Seg.	Temp.	Time	Cycle
1	95 °C	5min	
2	98°C	30s	20×
3	65 °C* (-0.5 °C/cycle)	30s	
4	72 °C	45s*	
5	98°C	30s	15×
6	55 °C*	30s	
7	72 °C	45s*	
8	72 °C	5min	
9	10 °C	hold	
PCR program II (the second choice)			
Seg.	Temp.	Time	Cycle
1	95 °C	5min	
2	98°C	30s	35×
3	58 °C*	30s	
4	72 °C	45s*	
5	72 °C	5min	
6	10 °C	hold	

Note\*: Annealing temperature and extension time can be determined according to the actual amplification situation and amplification enzyme efficiency.