


## Genotyping Report

|           |             |             |                 |                    |             |
|-----------|-------------|-------------|-----------------|--------------------|-------------|
| Strain ID | T038701     | Strain Type | CKO(Cas9)       | Genetic Background | C57BL/6JGpt |
| Designer  | Binjie Jiao | Gene Name   | <i>Preliid1</i> |                    |             |

### 1. Strategy of Genotyping




Legend:  Loxp    Primer   ① ② PCR Reaction Number    Sequencing Region

Wild type: ①PCR reaction obtains a single WT band; ②PCR reaction obtains a single WT band.  
Heterozygote: ①PCR reaction obtains a WT band and a Targeted band; ②PCR reaction obtains a WT band and a Targeted band.  
Homozygote: ①PCR reaction obtains a single Targeted band; ②PCR reaction obtains a single Targeted band.  
Note: The sizes of WT and Targeted band are shown below.

### 2. Primer Information

| PCR No.  | Primer No. | Primer Name    | Sequence                | Band Size                    |
|----------|------------|----------------|-------------------------|------------------------------|
| ①(5'arm) | F1         | T038701(P1)-F1 | GAATCAGTCCAAGCCTCCAAAG  | WT: 242bp<br>Targeted: 347bp |
|          | R1         | T038701(P1)-R1 | CCGACATGGATTCTGTGGCTC   |                              |
| ②(3'arm) | F2         | T038701(P1)-F2 | GTGCATGAGATTTGAGGAAGTGC | WT: 436bp<br>Targeted: 542bp |
|          | R2         | T038701(P1)-R2 | GAGCAGTGTTCCTCGTCTGCCT  |                              |

### 3. Gel Image & Conclusion



Note: P: Heterozygous samples; WT: Wildtype control; B: Blank control (ddH<sub>2</sub>O); M: DNA Ladder  
① Control (WT) : It is an important reference mark for whether the PCR reaction is successful and whether the

product band position and size meet the theoretical requirements.

② Control (B) : PCR amplification was performed without template in the PCR reagent to monitor whether the reagent was contaminated.

#### 4. PCR Condition

(Generally recommend to use Vazyme P222; If the sequences contain special structures such as GC%  $\geq$  60% or GC%  $\leq$  40%, recommend to use Vazyme P515.)

| PCR Reaction Component           |                                                                                        |      |             |
|----------------------------------|----------------------------------------------------------------------------------------|------|-------------|
| Seg.                             | reaction component                                                                     |      | Volume (μl) |
| 1                                | 2 × Rapid Taq Master Mix(Vazyme P222)<br>or<br>2 × Phanta Max Master Mix (Vazyme P515) |      | 12.5        |
| 2                                | ddH2O                                                                                  |      | 9.5         |
| 3                                | Primer A(10pmol/μl)                                                                    |      | 1           |
| 4                                | Primer B(10pmol/μl)                                                                    |      | 1           |
| 5                                | Template(20~80ng/μl)                                                                   |      | 1           |
| PCR program I priority selection |                                                                                        |      |             |
| Seg.                             | Temp.                                                                                  | Time | Cycle       |
| 1                                | 95℃                                                                                    | 5min | 20×         |
| 2                                | 98℃                                                                                    | 30s  |             |
| 3                                | 65℃* (-0.5℃/cycle)                                                                     | 30s  |             |
| 4                                | 72℃                                                                                    | 45s* |             |
| 5                                | 98℃                                                                                    | 30s  | 15×         |
| 6                                | 55℃*                                                                                   | 30s  |             |
| 7                                | 72℃                                                                                    | 45s* |             |
| 8                                | 72℃                                                                                    | 5min |             |
| 9                                | 10℃                                                                                    | hold |             |
| PCR program II the second choice |                                                                                        |      |             |
| Seg.                             | Temp.                                                                                  | Time | Cycle       |
| 1                                | 95℃                                                                                    | 5min | 35×         |
| 2                                | 98℃                                                                                    | 30s  |             |
| 3                                | 58℃*                                                                                   | 30s  |             |
| 4                                | 72℃                                                                                    | 45s* |             |
| 5                                | 72℃                                                                                    | 5min |             |
| 6                                | 10℃                                                                                    | hold |             |

Note\*: Annealing temperature and extension time can be determined according to the actual amplification situation



and amplification enzyme efficiency.