

Traf7 Cas9-KO Strategy

Designer: Qian Chen

Reviewer: Xiangli Bian

Design Date: 2023-12-19

Overview

Target Gene Name

• Traf7

Project Type


• Cas9-KO

Genetic Background

• C57BL/6JGpt



Strain Strategy



Technical Information

- The *Traf7* gene has 26 transcripts. According to the structure of *Traf7* gene, exon 6-exon 9 of *Traf7*-215 (ENSMUST00000176652.8) transcript is recommended as the knockout region. The region contains 446bp coding sequence. Knocking out the region will result in disruption of protein function.
- In this project we use CRISPR-Cas9 technology to modify *Traf7* gene. The brief process is as follows: gRNAs were transcribed in vitro. Cas9 and gRNAs were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and on-target amplicon sequencing. A stable F1-generation mouse strain was obtained by mating positive F0-generation mice with C57BL/6JGpt mice and confirmation of the desired mutant allele was carried out by PCR and on-target amplicon sequencing.



Gene Information

Traf7 TNF receptor-associated factor 7 [Mus musculus (house mouse)]

≛ Download Datasets

☆ ?

Gene ID: 224619, updated on 23-Nov-2023



Official Symbol Traf7 provided by MGI

Official Full Name TNF receptor-associated factor 7 provided by MGI

Primary source MGI:MGI:3042141

See related Ensembl: ENSMUSG00000052752 AllianceGenome: MGI: 3042141

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Muridae; Mus; Mus

Also known as RFWD1

Summary Predicted to enable ubiquitin-protein transferase activity. Acts upstream of or within several processes, including cellular response to interleukin-6; negative regulation of transcription by RNA

polymerase II; and positive regulation of protein sumoylation. Located in cytosol; nucleus; and perinuclear region of cytoplasm. Is expressed in genitourinary system and pancreas epithelium.

Orthologous to human TRAF7 (TNF receptor associated factor 7). [provided by Alliance of Genome Resources, Apr 2022]

Expression Ubiquitous expression in ovary adult (RPKM 69.0), lung adult (RPKM 60.1) and 28 other tissues See more

Orthologs <u>human</u> all

Try the new Gene table

Try the new Transcript table

Source: https://www.ncbi.nlm.nih.gov/




Transcript Information

The gene has 26 transcripts, all transcripts are shown below:

Transcript ID	Name	bp 🌲	Protein ▼	Biotype	CCDS	UniProt Match	Flags		
ENSMUST00000176652.8	Traf7-215	3003	<u>669aa</u>	Protein coding	CCDS50014 €	F8WJF7₁©	Ensembl Canonical GE	NCODE basic APPRIS P1	TSL:5
ENSMUST00000088464.12	Traf7-202	2504	669aa	Protein coding	CCDS50014 @	F8WJF7₫	GENCODE ba	sic APPRIS P1 TSL:5	
ENSMUST00000176353.8	Traf7-210	2503	629aa	Protein coding	CCDS50015 €	Q8C2K8₽	GENCODE basic TSL:1		
ENSMUST00000070777.13	Traf7-201	2384	629aa	Protein coding	CCDS50015 ₽	Q8C2K8₽	GENCO	ODE basic TSL:5	
ENSMUST00000176237.8	Traf7-208	950	271aa	Protein coding		H3BJD9₽	TSL:5	CDS 3' incomplete	
ENSMUST00000177154.8	Traf7-222	758	228aa	Protein coding		H3BLQ3₽	TSL:5	CDS 3' incomplete	
ENSMUST00000177193.8	Traf7-223	644	170aa	Protein coding		H3BK82₽	TSL:3	CDS 3' incomplete	
ENSMUST00000176086.8	Traf7-205	2687	<u>132aa</u>	Nonsense mediated decay		H3BLM6₽		TSL:1	
ENSMUST00000175698.8	Traf7-203	755	84aa	Nonsense mediated decay		H3BKN1₽	TSL:3	CDS 5' incomplete	
ENSMUST00000176668.8	Traf7-216	2667	81aa	Nonsense mediated decay		H3BKZ1₽		TSL:1	
ENSMUST00000177405.8	Traf7-225	606	80aa	Nonsense mediated decay		H3BJU4e₽		TSL:3	
ENSMUST00000177502.8	Traf7-226	641	76aa	Nonsense mediated decay		H3BJF9₽	TSL:3	CDS 5' incomplete	
ENSMUST00000176178.8	Traf7-207	680	54aa	Nonsense mediated decay		H3BJ15 €		TSL:3	
ENSMUST00000177401.8	Traf7-224	699	45aa	Nonsense mediated decay		H3BKC9₽		TSL:2	
ENSMUST00000176324.8	Traf7-209	604	44aa	Nonsense mediated decay		H3BKC7₽		TSL:3	
ENSMUST00000177025.2	Traf7-220	428	44aa	Nonsense mediated decay		H3BKC7@		TSL:3	
ENSMUST00000176092.2	Traf7-206	382	No protein	Protein coding CDS not defined		12		TSL:3	
ENSMUST00000177139.8	Traf7-221	332	No protein	Protein coding CDS not defined		-		TSL:3	
ENSMUST00000177024.8	Traf7-219	3194	No protein	Retained intron		-		TSL:1	
ENSMUST00000176805.2	Traf7-217	1749	No protein	Retained intron		100		TSL:5	
ENSMUST00000176530.8	Traf7-213	938	No protein	Retained intron		-		TSL:3	
ENSMUST00000176900.8	Traf7-218	861	No protein	Retained intron		(4)		TSL:3	
ENSMUST00000176361.2	Traf7-211	693	No protein	Retained intron		-		TSL:3	
ENSMUST00000176434.8	Traf7-212	559	No protein	Retained intron				TSL:2	
ENSMUST00000176633.2	Traf7-214	500	No protein	Retained intron		15.0		TSL:2	
ENSMUST00000175732.2	Traf7-204	366	No protein	Retained intron		(4)		TSL:1	


The strategy is based on the design of *Traf7*-215 transcript, the transcription is shown below:



Source: https://www.ensembl.org




Genomic Information





Source: : https://www.ensembl.org


Protein Information





Source: : https://www.ensembl.org

Mouse Phenotype Information (MGI)





Important Information

- According to MGI, Traf7 knockout homozygous mice died.
- *Traf7* is located on Chr17. If the knockout mice are crossed with other mouse strains to obtain double homozygous mutant offspring, please avoid the situation that the second gene is on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risks of the mutation on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

