


Genotyping Report

Strain ID	T052691	Strain Type	TG	Genetic Background	C57BL/6JGpt
Designer	Tianjiao Wang	Gene Name	Sox2-iCre		


1. Strategy of Genotyping



2. Primer Information

PCR No.	Primer No.	Primer No.	Sequence	Band Size
① 5'connector	F1	T052691-F1	ACTTCCTTTGTCCCAAATCTGTGC	WT:0bp Targeted:801bp
	R1	T052691-R1	CTGATGTCCTGGCATCTGTCAGAGTT	
② 3'connector	F2	T052691-F2	CAAGGATGACTCTGGGCAGAG	WT:0bp Targeted:350bp
	R2	T052691-R2	AGCCAGAAGTCAGATGCTCAAG	

3. Gel Image & Conclusion



product band position and size meet the theoretical requirements.

② Control (B) : PCR amplification was performed without template in the PCR reagent to monitor whether the reagent was contaminated.

4. PCR Condition

(Generally recommend to use Vazyme P222; If the sequences contain special structures such as GC% $\geq$ 60% or GC% $\leq$ 40%, recommend to use Vazyme P515.)

PCR Reaction Component			
Seg.	reaction component		Volume (μl)
1	2 × Rapid Taq Master Mix(Vazyme P222) or 2 × Phanta Max Master Mix (Vazyme P515)		12.5
2	ddH2O		9.5
3	Primer A(10pmol/μl)		1
4	Primer B(10pmol/μl)		1
5	Template(20~80ng/μl)		1
PCR program I priority selection			
Seg.	Temp.	Time	Cycle
1	95℃	5min	20×
2	98℃	30s	
3	65℃* (-0.5℃/cycle)	30s	
4	72℃	45s*	
5	98℃	30s	15×
6	55℃*	30s	
7	72℃	45s*	
8	72℃	5min	
9	10℃	hold	
PCR program II the second choice			
Seg.	Temp.	Time	Cycle
1	95℃	5min	35×
2	98℃	30s	
3	58℃*	30s	
4	72℃	45s*	
5	72℃	5min	
6	10℃	hold	

Note*: Annealing temperature and extension time can be determined according to the actual amplification

situation and amplification enzyme efficiency.